CO₂ Laser-induced Endotracheal Fire

Jae-Gu Cho
Min Woo Park
Seung-Kuk Baek
Soon-Young Kwon
Kwang-Yoon Jung
Jeong-Soo Woo

*Department of Otolaryngology-Head and Neck
Surgery, Korea University College of Medicine,
Seoul, Korea*

The objective of this paper is to describe long-term complications and their management in contrast to acute measures after endotracheal laser-induced fire. This case pertains to a 56-year-old patient in whom an endotracheal fire occurred during CO₂ laser surgery. Despite local swelling and evidence of acute lung injury, the patient was extubated the following day under single-shot cortisone and inhalation of dispersed adrenaline under assisted spontaneous breathing. Regular flexible bronchoscopy and spirometry were used to assess wound healing. But 14 weeks after uneventful recovery, the patient presented with an acute inspiratory stridor, related to a tracheal stenosis 2.5 cm distal to the glottic level. Further follow-up was uneventful after the detection of tracheal end-to-end anastomosis. Early extubation under ITU conditions avoided the need for tracheostomy and its sequelae. However, tracheal stenosis did not become apparent before week 14. While a conservative approach was successfully implemented in the acute management of laser-induced endotracheal fire, the risk of further long-term complications implies the need for a prolonged follow-up regime even in cases of less extensive burns.

Key words

Laser; Burn; Trachea

Received June 10, 2013
Revised June 11, 2013
Accepted June 11, 2013

Correspondence

Jae-Gu Cho
Department of Otorhinolaryngology-Head and
Neck Surgery, Guro Hospital, Korea University
College of Medicine, 80, Guro-dong, Guro-gu,
Seoul 152-703, Korea
Tel: +82-2-2626-3189
Fax: +82-2-868-0475
E-mail: jgcho@korea.ac.kr

© Korean Society for Laser Medicine and Surgery

© This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

The objective of this presentation is to outline long-term complications and their management in contrast to acute measures after endotracheal laser-induced fire. This case focuses on a 56-year-old patient in whom an endotracheal fire occurred during CO₂ laser surgery. Despite local swelling and evidence of acute lung injury, the patient was extubated the following day under single-shot cortisone and inhalation of dispersed adrenaline under assisted spontaneous breathing. Wound healing was assessed by regular flexible bronchoscopy and spirometry. Fourteen weeks after uneventful recovery, the patient presented with acute inspiratory stridor, related to a tracheal stenosis 2.5 cm distal to the glottic level. After tracheal end-to-end anastomosis, further follow-up was uneventful. Early extubation under ITU conditions avoided the need for tracheostomy and its sequelae. However, tracheal stenosis did not become apparent before week 14. While in acute management of laser-induced endotracheal fire a conservative approach was established successfully, the risk of further longterm complications implies the need for a prolonged follow-up regime even in cases of less extensive burns.

CASE REPORT

A 56-year-old patient was referred to the anaesthesiology intensive care unit from a district general hospital on 23 December 1999. Six hours prior to the referral, this patient had undergone excision of bilateral vocal fold granulomas with a CO₂ laser under general anaesthesia when ignition of the endotracheal tube followed by combustion of inflammable gases occurred. A metal-sheathed laser-resistant endotracheal tube had been used. The O₂-concentration was kept at 30 per cent maximum while intravenous anaesthetics were employed. Laser power and mode were unknown.

On arrival we saw an intubated and sedated patient. Both ears, nasal cavities and the nasopharynx were unremarkable. The oropharynx, in particular the base of the tongue showed marked inflammation with traces of carbonized material. Both aryepiglottic folds and the epiglottis were markedly swollen and covered with fibrinous exudate. The O₂-saturation was kept between 93 to 96 per cent under 30 per cent oxygen delivered by pressure-controlled ventilation with positive end-expiratory pressure. Maximum airway pressure was limited to 30 mmHg. All other vital parameters were within normal limits. No cardiocirculatory support was

Fig. 1. Day 21 post-incidence. Marked fibrinous coating of both vocal folds, subglottic space.

needed.

On day 10 the patient was referred to the Otorhinolaryngology ward from the intensive care unit. Minimal invasive ventilation with 41/minute oxygen was continued during five periods of 10 minutes each during daytime. Although oxygen concentration occasionally dropped below 92 per cent, the patient did not complain of any further shortness of breath. Mild infiltration, markedly in the right superior lobe, had almost disappeared on the chest X-ray by day 12. Spirometry showed normal results, and the patient was discharged from hospital on day 16 (Fig. 1).

During out-patient follow-up, hoarseness continued and full re-epithelialization of both vocal folds was complete by week 10. Spirometry showed mild obstruction only. A ventilation/perfusion scintigram confirmed the finding of a mild ventilation deficit in the right upper lobe, otherwise there were no further abnormalities.

During week 14 the patient presented with acute inspiratory stridor. While clinical examination showed only minimal synechia at the vocal fold level, a coronal reconstruction of an axial computed tomography (CT) scan revealed a tracheal stenosis, about 2.5 cm distal to the glottic level, reducing the tracheal diameter to 7.5 mm (Fig. 2, 3). Functionally, this resulted in severe irreversible respiratory obstruction. Thus, partial tracheal resection with an end-to-end anastomosis was performed the following week. The patient was extubated under bronchoscopy three days later with no further dyspnoea.

A follow-up bronchoscopy under general anaesthesia during week 19 showed some granulating tissue in

Fig. 2. Pre-operative CT scan in sagittal plane reconstruction with tracheal stenosis. X, glottis level; T, thyroid cartilage.

the anastomosis region, that could be removed with a Nd:YAG laser at 10 W in contact mode. Further respiratory function tests were all normal. The patient, who is currently undergoing voice therapy, complains of some episodes of shortness of breath while speaking. This could be related to the remaining mucosal defect on the vocal fold level. Otherwise, he is back at work with no further complaints.

DISCUSSION

Following the increasing experience with laser-tissue interaction, catastrophic events such as endotracheal fires have become more and more rare. While training in prevention is important, the authors present this case in order to outline a possible low-interventional regime in the unfortunate case when an endotracheal fire occurs. In current literature, suggestions are made to avoid laser-induced fires by either employing jet ventilation or ventilation via a rigid bronchoscope.^{1,2} However, some authors do admit that the risk of endotracheal fire cannot be eliminated completely, especially if endotracheal tubes are employed.¹⁻⁴ Emergency procedures comprise immediately stopping the oxygen and anaesthetic gas flow, the removal of burning material, extinguishing the fire and reintubation with a endotracheal tube followed by immediate bronchoscopy.^{1,3,5,6} Yet there is little reported about the sub-acute and long-term management of an endotracheal fire. Ossoff et al. noted that most severe burns were seen after polyvinyl chloride tube ignition in an animal experiment, but warned that ash from silicone

Fig. 3. Rigid bronchoscopic evaluation after 14 weeks post-incidence.

tubes could induce silicosis as a long-term effect.

Schramm et al.⁵ suggested frequent rigid bronchoscopy and early tracheostomy in order to aid ventilation under positive end-expiratory pressure. While we followed this regime in general, we decided to extubate the patient as early as possible and introduced continuous positive airway pressure ventilation under O₂ saturation monitoring instead. The rationale was 1) to avoid tracheostomy which could be a cause for additional inflammation and possible tracheal wall necrosis and 2) to allow mucociliary clearance of the upper airways at an early stage without additional irritation from the exterior. The authors regard this management as feasible as long as there is no severe laryngeal or pharyngeal oedema, entailing the need for tracheostomy, or severe pulmonary oxygenation deficit. To answer the latter, a ventilation/perfusion scintigram proved to be helpful. This case demonstrates that a conservative approach in the acute phase management can help to avoid secondary complications during this period - however, a marked tracheal stenosis did not become clinically apparent before week 14. Therefore, we recommend follow-up of these patients for about six months. Most information was gained from pulmonary function tests in conjunction with a coronary reconstruction of an axial neck CT scan, that eventually made the case for tracheal end-to-end anastomosis.

REFERENCES

1. Hermens JM, Bennett MJ, Hirshman CA. Anesthesia for laser surgery. *Anesth Analg* 1983;62:218-29.

2. Hunsaker DH. Anesthesia for microlaryngeal surgery: the case for subglottic jet ventilation. *Laryngoscope* 1994;104(8 Pt 2 Suppl 65):1-30.
3. Ossoff RH, Duncavage JA, Eisenman TS, Karlan MS. Comparison of tracheal damage from laser-ignited endotracheal tube fires. *Ann Otol Rhinol Laryngol* 1983;92:333-6.
4. Airway fires: reducing the risk during laser surgery. *Health Devices* 1990;19:109-39.
5. Schramm VL Jr, Mattox DE, Stool SE. Acute management of laser-ignited intratracheal explosion. *Laryngoscope* 1981;91:1417-26.
6. Bingham HG, Gallagher TJ, Singleton GT, Gravenstein JS, Pashayan AG, Bjoraker DG. Carbon dioxide laser burn of laryngotracheobronchial mucosa. *J Burn Care Rehabil* 1990;11:64-6.